Statistical properties of the entropy from ordinal patterns

Cite as: Chaos **32**, 113118 (2022); doi: 10.1063/5.0118706 Submitted: 5 August 2022 · Accepted: 22 September 2022 · Published Online: 4 November 2022



E. T. C. Chagas,^{1,a)} (b) A. C. Frery,² (b) J. Gambini,³ M. M. Lucini,^{4,5} H. S. Ramos,¹ and A. A. Rey⁶

AFFILIATIONS

¹Departamento de Ciência da Computação, Universidade Federal de Minas Gerais, Belo Horizonte, MG 30123-970, Brazil ²School of Mathematics and Statistics, Victoria University of Wellington, Wellington 6140, New Zealand ³Departamento de Ingeniería Informática, Instituto Tecnológico de Buenos Aires, Av. Madero 399, Buenos Aires C1106ACD, Argentina

⁴Facultad de Ciencias Exactas, Naturales y Agrimensura, Universidad Nacional de Nordeste, Av. Libertad 5450—Campus "Deodoro Roca," 3400 Corrientes, Argentina

⁵Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina (CONICET), Godoy Cruz 2290, Buenos Aires, Argentina
⁶Centro de Procesamiento de Se nales e Imágenes, Department of Mathematics, Universidad Tecnológica Nacional Facultad Regional Buenos Aires, Ciudad de Buenos Aires C1179AAQ, Argentina

Note: This paper is part of the Focus Issue on Ordinal Methods: Concepts, Applications, New Developments and Challenges. ^{a)}Author to whom correspondence should be addressed: eduarda.chagas@dcc.ufmg.br

ABSTRACT

The ultimate purpose of the statistical analysis of ordinal patterns is to characterize the distribution of the features they induce. In particular, knowing the joint distribution of the pair entropy-statistical complexity for a large class of time series models would allow statistical tests that are unavailable to date. Working in this direction, we characterize the asymptotic distribution of the empirical Shannon's entropy for any model under which the true normalized entropy is neither zero nor one. We obtain the asymptotic distribution from the central limit theorem (assuming large time series), the multivariate delta method, and a third-order correction of its mean value. We discuss the applicability of other results (exact, first-, and second-order corrections) regarding their accuracy and numerical stability. Within a general framework for building test statistics about Shannon's entropy, we present a bilateral test that verifies if there is enough evidence to reject the hypothesis that two signals produce ordinal patterns with the same Shannon's entropy. We applied this bilateral test to the daily maximum temperature time series from three cities (Dublin, Edinburgh, and Miami) and obtained sensible results.

Published under an exclusive license by AIP Publishing. https://doi.org/10.1063/5.0118706

We present results about the statistical properties of ordinal patterns' Shannon entropy: exact and approximate first-, second-, and third-order moments and its asymptotic distribution. We discuss their limitations and propose a new model. We present applications to test the hypothesis that two time series produce ordinal patterns with the same Shannon entropy.

I. INTRODUCTION

The analysis of signals throughout their ordinal patterns has received much attention since it was proposed by Bandt and Pompe.^{\perp} This approach is appealing, among other virtues, for its ability to translate analysis into a graphical depiction: the signal is

represented as a point in the entropy–complexity plane ($H \times C$), a closed two-dimensional manifold.

There is a vast literature of successful applications of this kind of analysis, among them: The ability to distinguish noise from chaos,² the analysis of economic markets and biomedical systems,³ fault diagnosis of rotating machinery,⁴ and image texture analysis.⁵⁻⁷

Chagas *et al.*⁸ pointed out a common trait of such applications: the lack of a theoretical background when performing comparisons among signals and when contrasting a signal with a hypothesized model. The authors proposed a test for the white noise model using empirical evidence. Although this approach proved useful, extending it to other situations requires extensive simulation experiments and data analysis.

The ultimate goal in a statistical approach to the Bandt and Pompe methodology is the exact specification of the distribution of the coordinates of points in $H \times C$ for a large class of time series models. Our purpose is delimited: In this work, we present exact, approximate, and asymptotic results of the distribution of Shannon's entropy when the patterns that result from the Bandt and Pompe symbolization follow a multinomial distribution. The properties of such a distribution that we employ later are recalled in Sec. II.

The exact and approximate results we present in Sec. III appeared in the works by Basharin,⁹ by Hutcheson,¹⁰ and by Hutcheson and Shenton.¹¹ These works have received little attention from the community that uses the permutation entropy approach.

We discuss the applicability of these expressions in Sec. IV, stressing that the exact ones are of limited use in practice with highstandard numerical platforms currently available. We obtain the asymptotic distribution of Shannon's entropy using the properties of the multinomial distribution, the central limit theorem, and the multivariate delta method (Sec. III D). We propose a better approximation by plugging a third-order correction of the mean into its asymptotic expression [Eq. (29)].

We also provide a general framework for hypotheses testing, including the possibility of contrasting series of different lengths and embedding dimensions. We explicitly derive test statistics for the null hypothesis that two time series produce the same number of symbols, and we apply these tests to climatology data.

II. ORDINAL PATTERNS AND THE MULTINOMIAL DISTRIBUTION

Let $\mathbf{x} = (x_1, x_2, ..., x_{n+D-1})$ be a real-valued time series of length n + D - 1 without ties. We compute $\pi_1, \pi_2, ..., \pi_n$ symbols from subsequences of embedding dimension *D*, for instance, $\pi_j = BP(x_j, x_{j+1}, ..., x_{j+D-1})$, where BP is the Bandt–Pompe symbolization. There are *D*! possible symbols: $\pi_j \in \mathbf{\pi} = \{\pi^1, \pi^2, ..., \pi^{D!}\}$. We form the histogram of proportions $\mathbf{h} = (h_1, h_2, ..., h_{D!})$ in which the bin h_ℓ is the proportion of symbols of type π^ℓ of the total *n* symbols. We will model those symbols as a *k* dimensional random vector with a multinomial distribution in which k = D!.

Consider a series of *n* independent trials in which only one of *k* mutually exclusive events $\pi^1, \pi^2, \ldots, \pi^k$ is observed with probability p_1, p_2, \ldots, p_k , respectively, such that $p_\ell \ge 0$ and $\sum_{\ell=1}^k p_\ell = 1$. Let $N = (N_1, N_2, \ldots, N_k)$ be the vector of random variables that count the number of occurrences of the events $\pi^1, \pi^2, \ldots, \pi^k$ in the *n* trials, with $\sum_{\ell=1}^k N_\ell = n$. Then, the joint distribution of *N* is

$$\Pr\left(\mathbf{N} = (n_1, n_2, \dots, n_k)\right) = n! \prod_{\ell=1}^k \frac{p_{\ell}^{n_\ell}}{n_{\ell}!},$$
 (1)

where $n_{\ell} \geq 0$ and $\sum_{\ell=1}^{k} n_{\ell} = n$. We denote this situation as $N \sim \text{Mult}(n, p)$, with $p = (p_1, p_2, \dots, p_k)$.

As per the conditions under which the Bandt and Pompe technique is used, we require that $n \gg k$. The common practice is k = 3!, 4!, 5!, or 6!, and $n \ge 100k$.

Consider the random vector $N \sim Mult(n, p)$. Its main moments are

$$\mathbf{E}(N_{\ell}) = np_{\ell},\tag{2}$$

$$\operatorname{Var}(N_{\ell}) = np_{\ell}(1 - p_{\ell}), \tag{3}$$

$$\operatorname{Cov}(N_{\ell}, N_j) = -np_{\ell}p_j, \text{ and }$$
(4)

$$\rho(N_{\ell}, N_j) = \sqrt{\frac{p_{\ell} p_j}{(1 - p_{\ell})(1 - p_j)}}$$
(5)

for every $1 \le \ell, j \le k$. The book by Johnson *et al.* (Sec. 11.18 in Ref. 12) is a comprehensive treatise on this topic.

The maximum likelihood (ML) estimator of p_{ℓ} is the relative frequency $\hat{p}_{\ell} = N_{\ell}/n$, $1 \le \ell \le k$, and the distribution of $n\hat{p}$ is Mult(n, p). The properties of ML estimators grant that if \hat{p} is the ML estimator of p, then for any function g(p), the ML estimator of g(p), namely, $\hat{g}(p)$, is $g(\hat{p})$; cf. Casella and Berger (Theorem 7.2.10 in Ref. 13). We will use this result to obtain the asymptotic distribution of Shannon's entropy.

Let $X_n = (X_{1n}, X_{2n}, \dots, X_{kn})$ be a sequence of independent and identically distributed random vectors, with a Mult(n, p) distribution. If \hat{p} is the vector of sample proportions and $Y_n = \sqrt{n}(\hat{p} - p)$, then

$$\mathbf{E}(\boldsymbol{Y}_n) = \mathbf{0},\tag{6}$$

$$Cov(\boldsymbol{Y}_n) = \mathbf{D}_{\boldsymbol{p}} - \boldsymbol{p}\boldsymbol{p}^{\mathrm{T}},\tag{7}$$

where $\mathbf{D}_p = \text{Diag}(p_1, p_2, \dots, p_k)$, and the superscript T denotes transposition. Mukhopadhyay (Theorem A.2.1 in Ref. 14) states that

$$Y_n \xrightarrow{\mathscr{D}} \mathscr{N}(\mathbf{0}, \mathbf{D}_p - pp^{\mathrm{T}}).$$
(8)

This asymptotic distribution is the basis of our forthcoming derivations.

III. THE ENTROPY AND ITS PROPERTIES

This section presents the main results about $H(\hat{p})$ under the multinomial model. Regarding its mean and variance, we recall exact expressions, first, second, and third-order approximations. Finally, its asymptotic distribution is presented.

The Shannon's entropy of a multinomial-distributed random variable is

$$H(\boldsymbol{p}) = -\sum_{\ell=1}^{k} p_{\ell} \ln p_{\ell}, \qquad (9)$$

which is bounded between 0 and ln k. The minimum is attained when $p_{\ell} = 1$ for some $1 \le l \le k$ and $p_j = 0$ for every $j \ne \ell$, while the expression is maximized by $p_{\ell} = 1/k$ for every $1 \le \ell \le k$. In the following, we will consider only probability vectors *p* that differ from these two extreme points.

We are interested in the statistical properties of $H(\mathbf{p})$ when it is indexed by $\hat{\mathbf{p}} = (\hat{p}_1, \hat{p}_2, \dots, \hat{p}_k)$, the ML estimator of \mathbf{p} . Our problem then becomes in finding the distribution of

(11)

$$H(\hat{p}) = -\sum_{\ell=1}^{k} \hat{p}_{\ell} \ln \hat{p}_{\ell}$$
(10)
$$= -\sum_{\ell=1}^{k} \frac{N_{\ell}}{n} \ln \frac{N_{\ell}}{n}$$
$$= \ln n - \frac{1}{n} \sum_{\ell=1}^{k} N_{\ell} \ln N_{\ell}$$
(11)

under $N = (N_1, N_2, \ldots, N_k) \sim \text{Mult}(n, p)$.

A. Exact mean and variance

Hutcheson and Shenton¹¹ found the expressions for the exact first and second-order moments of (10). They are given by

$$E(H(\hat{\boldsymbol{p}})) = \ln n - \sum_{j=1}^{n-1} \binom{n-1}{n-j} \ln(n-j+1) \sum_{\ell=1}^{k} p_{\ell}^{n-j+1} (1-p_{\ell})^{j-1},$$
(12)

and, denoting the integer part of $x \in \mathbb{R}$ as [x],

$$\operatorname{Var}(H(\widehat{p})) = \sum_{a=0}^{n-2} \binom{n-1}{a} \sum_{\ell=1}^{k} p_{\ell}^{n-a} (1-p_{\ell})^{a} \left[\sum_{b=a+1}^{n-1} \binom{n-1}{b} \sum_{\ell=1}^{k} p_{\ell}^{n-b} (1-p_{\ell})^{b} \left(\ln \frac{n-a}{n-b} \right)^{2} \right] - \frac{n-1}{n} \sum_{b=0}^{n-3} \binom{n-2}{b} \left[\sum_{a=0}^{\left[(n-b-2)/2\right]} \binom{n-b-2}{a} \sum_{\ell \neq j} p_{\ell}^{n-a-b-1} (1-p_{\ell})^{a+1} (1-p_{\ell}p_{j})^{b} \left(\ln \frac{n-a-b-1}{a+1} \right)^{2} \right].$$
(13)

B. First-order approximation

Basharin⁹ provided some of the first results about the properties of $H(\hat{p})$. This author found first-order approximations for its expected value and variance, whose expressions are given by

$$\check{\mathrm{E}}\big(H(\hat{\boldsymbol{p}})\big) = H(\boldsymbol{p}) - \frac{k-1}{2n\ln 2}$$
(14)

and

$$\check{\operatorname{Var}}(H(\hat{\boldsymbol{p}})) = \frac{1}{n} \Big[\sum_{\ell=1}^{k} p_{\ell} \frac{\ln^2 p_{\ell}}{\ln^2 2} - H^2(\boldsymbol{p}) \Big].$$
(15)

Notice that Eq. (14) can be used to build a first-order bias corrected estimator for $H(\mathbf{p})$. Basharin⁹ also proved that $H(\hat{\mathbf{p}})$ is consistent $[H(\hat{p}) \rightarrow H(p) \text{ when } n \rightarrow \infty]$ and asymptotically normal.

C. Third-order approximation

Hutcheson¹⁰ presented third-order approximate expressions by expanding (12) and (13) in series and retaining only the initial terms. With this, we have

$$\tilde{E}(H(\hat{p})) = H(p) - \frac{k-1}{2n} + \frac{1 - \sum_{\ell=1}^{k} p_{\ell}^{-1}}{12n^2} + \frac{\sum_{\ell=1}^{k} (p_{\ell}^{-1} - p_{\ell}^{-2})}{12n^3}$$
(16)

and

$$\widetilde{\operatorname{Var}}(H(\widehat{\boldsymbol{p}})) = \frac{\sum_{\ell=1}^{k} p_{\ell} \ln^{2} p_{\ell} - \left(\sum_{\ell=1}^{k} p_{\ell} \ln p_{\ell}\right)^{2}}{n} + \frac{k-1}{2n^{2}} + \frac{\sum_{\ell=1}^{k} p_{\ell}^{-1} - \sum_{\ell=1}^{k} p_{\ell}^{-1} \ln p_{\ell} + \sum_{\ell=1}^{k} p_{\ell}^{-1} \sum_{\ell=1}^{k} p_{\ell} \ln p_{\ell} - 1}{6n^{3}}.$$
(17)

Notice that dropping the last term in these expressions yields second-order approximations for the mean and variance, respectively. Exact expressions should be better than approximations, but we will see in Sec. IV that the first ones have limited application due to the numerical instabilities they incur. On the contrary, all approximations are numerically stable.

Hutcheson and Shenton¹¹ briefly discussed how to obtain higher-order moments that would be useful for computing the skewness and kurtosis.

ARTICLE

D. Asymptotic distribution

We recall the following theorems known as the Delta Method and its multivariate version. For their proofs, we refer to Lehmann and Casella.¹⁵

Theorem 1. Let X_n be a sequence of independent and identically distributed random variables such that $\sqrt{n}(X_n - \theta)$ converges in distribution to a $\mathcal{N}(0, \sigma^2)$ law. Consider the transformation $h(X_n)$ such that $h'(\theta)$ exists and does not vanish. Then, $\sqrt{n}[h(X_n) - h(\theta)]$ converges in distribution to a $\mathcal{N}(0, \sigma^2[h'(\theta)]^2)$ law.

Theorem 2. Let $X_n = (X_{1n}, X_{2n}, ..., X_{kn})$ be a sequence of independent and identically distributed vectors of random variables such that $\sqrt{n}(X_{1n} - \theta_1, X_{2n} - \theta_2, ..., X_{kn} - \theta_k)$ converges in distribution to the multivariate normal law $\mathcal{N}_n(\mathbf{0}, \Sigma)$, where Σ is the covariance matrix. Suppose that $h_1, h_2, ..., h_k$ are real-functions continuously differentiable in a neighborhood of the parameter point $\boldsymbol{\theta} = (\theta_1, \theta_2, ..., \theta_k)$ such that the matrix of partial derivatives $B = (\partial h_i / \partial \theta_j)_{ij=1}^k$ is non-singular in the mentioned neighborhood. Then, the following convergence in distribution holds:

$$\frac{\sqrt{n} [h_1(\mathbf{X}_n) - h_1(\boldsymbol{\theta}), h_2(\mathbf{X}_n) - h_2(\boldsymbol{\theta}), \dots, h_k(\mathbf{X}_n) - h_k(\boldsymbol{\theta})]}{\xrightarrow{\mathscr{D}} \mathcal{N} (\mathbf{0}, B \Sigma B^T).}$$

For our case of interest $N \sim Mult(n, p)$, the covariance matrix of Eq. (8) is

$$\left(\mathbf{D}_{\boldsymbol{p}} - \boldsymbol{p}\boldsymbol{p}^{\mathrm{T}}\right)_{\ell j} = \begin{cases} p_{\ell}(1 - p_{\ell}) & \text{if } \ell = j, \\ p_{\ell}p_{j} & \text{if } \ell \neq j \end{cases}$$
(18)

for $1 \leq l, j \leq k$.

In order to apply the delta method using Theorem 2 to Shannon's entropy defined in Eq. (10), we use the functions

$$h_{\ell}(p_1, p_2, \dots, p_k) = p_{\ell} \ln p_{\ell},$$
 (19)

which verify that

$$\frac{\partial h_{\ell}}{\partial p_{j}} = \begin{cases} \ln p_{\ell} + 1 & \text{if } \ell = j, \\ 0 & \text{otherwise} \end{cases}$$
(20)

for $1 \le \ell, j \le k$. Hence, the covariance matrix of the multivariate normal limit distribution $\Sigma_p = (\partial h_\ell / \partial p_j) (\mathbf{D}_p - p p^{\mathrm{T}}) (\partial h_\ell / \partial p_j)^{\mathrm{T}}$ is of the form

$$(\Sigma_p)_{\ell j} = \begin{cases} (p_{\ell} - p_{\ell}^2)(\ln p_{\ell} + 1)^2 & \text{if } \ell = j, \\ -p_{\ell} p_j(\ln p_{\ell} + 1)(\ln p_j + 1) & \text{otherwise} \end{cases}$$
(21)

for $1 \le l, j \le k$. Therefore, we conclude that

$$\sqrt{n} \Big[h_1(\hat{p}_1) - h_1(p_1), h_2(\hat{p}_2) - h_2(p_2), \dots, h_k \hat{p}_k - h_k(p_k) \Big]$$

$$\xrightarrow{\mathscr{D}} \mathscr{N}(\mathbf{0}, \Sigma_p).$$
(22)

An equivalent expression is

$$\sqrt{n} \Big[h_1(\hat{p}_1), h_2(\hat{p}_2), \dots, h_k(\hat{p}_k) \Big] \xrightarrow{\mathscr{D}} \mathscr{N} \left(\sqrt{n} \begin{pmatrix} h_1(p_1) \\ h_2(p_2) \\ \vdots \\ h_k(p_k) \end{pmatrix}, \Sigma_p \right).$$
(23)

For a random vector Y such that $\sqrt{n}Y \xrightarrow{\mathcal{D}} \mathcal{N}(\sqrt{n}\mu, \Sigma)$, it can be proved that $E(\sqrt{n}Y) \rightarrow \sqrt{n}\mu$ and that $Var(\sqrt{n}Y) \rightarrow \Sigma$. Provided well-known properties, it holds that E(Y) $\rightarrow \mu$ and $Var(Y) \rightarrow 1/n\Sigma$. Applying this to Eq. (23),

$$\left[h_1(\hat{p}_1), h_2(\hat{p}_2), \dots, h_k(\hat{p}_k)\right] \xrightarrow{\mathscr{D}} \mathscr{N}\left(\begin{pmatrix}h_1(p_1)\\h_2(p_2)\\\vdots\\h_k(p_k)\end{pmatrix}, \frac{1}{n}\Sigma_p\right). \quad (24)$$

We now use Eq. (24) and the fact that the Shannon's entropy is a linear combination of the functions $\{h_1(p_1), h_2(p_2), \ldots, h_k(p_k)\}$.

Let $\mathbf{Z} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$ be a multivariate Gaussian k dimensional vector, with $\boldsymbol{\mu} \in \mathbb{R}^k$ and $\Sigma = (\sigma_{\ell,j}), 1 \leq \ell, j \leq k$. Let $W = \boldsymbol{a}^T \mathbf{Z}$ be a linear combination of the \mathbf{Z} elements, with $\boldsymbol{a} \in \mathbb{R}^k$. Thus, W is $\mathcal{N}(\boldsymbol{a}^T\boldsymbol{\mu}, \sum_{\ell=1}^k a_\ell^2 \sigma_{\ell,\ell} + 2\sum_{\ell=1}^{k-1} \sum_{j=\ell+1}^k a_\ell a_j \sigma_{\ell,j})$ distributed (see Lehmann and Casella¹⁵). Using the limit distribution presented in Eq. (24) and $\boldsymbol{a} = (-1, -1, \dots, -1)$, we have

$$H(\hat{\boldsymbol{p}}) = -\sum_{\ell=1}^{k} \hat{p}_{\ell} \ln \hat{p}_{\ell} \xrightarrow{\mathscr{D}} \mathscr{N} (H(\boldsymbol{p}), \sigma_{n,\boldsymbol{p}}^{2}), \qquad (25)$$

where

$$\sigma_{n,p}^{2} = \frac{1}{n} \sum_{\ell=1}^{k} p_{\ell} (1 - p_{\ell}) (\ln p_{\ell} + 1)^{2} - \frac{2}{n} \sum_{j=1}^{k-1} \sum_{\ell=j+1}^{k} p_{j} p_{\ell} (\ln p_{j} + 1) (\ln p_{\ell} + 1).$$
(26)

Notice that the asymptotic mean coincides with Eq. (9) and that computing the asymptotic variance does not pose any numerical difficulty.

Some studies and applications use a normalized entropy computed from Eq. (9) divided by $\ln k$. In this case, the asymptotic distribution is $\mathcal{N}(H(\mathbf{p})/\ln k, \sigma_{n,\mathbf{p}}^2/(\ln k)^2)$.

IV. EXPERIMENTS, NUMERICAL STABILITY, AND ACCURACY

Basharin's first-order approximations to the mean and variance, Eqs. (14) and (15), do not offer numerical difficulty, except in situations where there is at least one probability value near zero, $p_{\ell} \approx 0$ for some $1 \le \ell \le k$. Such cases are easily handled as the limit $\lim_{p\to 0} p \ln p = 0$ and pose no numerical challenges. The same situation occurs with second-order approximations, Eqs. (16) and (17).

Hutcheson's exact expressions, Eqs. (12) and (13), involve more sources of numerical instabilities, namely, combinatorial numbers, $\binom{a}{b}$ with very large *a* and *b* values. We implemented these exact expressions using computer algebra platforms (Mathematica, Yacas, and Maxima) as well as with high-precision specialized numerical functions in R (the **VeryLargeIntegers** package, that allows storing and operating with arbitrarily large integers). None of these platforms returned useful values of either Eq. (12) or Eq. (13) for practical situations. Despite that and for the sake of completeness, we report some of those results in the following.



FIG. 1. Log-relative errors for Basharin, Hutcheson, and asymptotic approximations using different underlying distributions: [LRE] = 2 (green), [LRE] = 1 (blue), and [LRE] = 0 (red).

Following Almiron *et al.*,¹⁶ we compare a "certified" value *c* with its "approximation" x by computing the absolute value of the relative error and taking its decimal logarithm,

$$LRE(x,c) = \begin{cases} -\log \frac{|x-c|}{|c|} & \text{if } c \neq 0, \text{ and} \\ -\log |x| & \text{otherwise.} \end{cases}$$
(27)

This log-relative error relates to the number of significant digits that are correctly computed; therefore, we report its integer part [LRE(x, c)].

In the following, the Basharin, Hutcheson, and asymptotic approximations are compared with the exact expected value given in (12).

In this experiment, we applied four different types of underlying distributions: equiprobable \mathcal{P}_e , a perturbed equiprobable law



FIG. 2. Relative errors for Basharin (pink), Hutcheson (green), and asymptotic (light blue) approximations for a given number of ordinal patterns k and different underlying distributions \mathcal{P}_{e} , \mathcal{P}_{2} , \mathcal{P}_{H} , and \mathcal{P}_{L} .

 $\mathcal{P}_2,$ the half perturbed distribution $\mathcal{P}_H,$ and the linear distribution $\mathcal{P}_L.$

- $\mathcal{P}_e: p_\ell = \frac{1}{k}, \ell = 1, \ldots, k.$
- \mathcal{P}_2 : $p_\ell = 1/k$, $\ell = 1, \dots, k-2$, $p_{k-1} = \frac{1}{k} + \varepsilon$, and $p_k = \frac{1}{k} \varepsilon$, $0 < \varepsilon < 1/k$.
- \mathcal{P}_{H} : $p_{\ell} = 1/k \varepsilon$, $\ell = 1, \dots, \frac{k}{2}$, and $p_{\ell} = 1/k + \varepsilon$, $\ell = \frac{k}{2}, \dots, k$, $0 < \varepsilon < 1/k$.

•
$$\mathcal{P}_{\mathrm{L}}: p_{\ell} = \frac{\iota}{\sum_{i=1}^{k} j}.$$

The results obtained are shown in Fig. 1, where green, blue, and red boxes correspond to [LRE] = 2, 1, and 0, respectively. It can be seen that the asymptotic approximation has the best performance, followed by Basharin's approximation.

The same scenarios and underlying distributions were used to compute the relative error

$$\operatorname{RE}(x,c) = \begin{cases} \frac{|x-c|}{|c|} & \text{if } c \neq 0, \text{ and} \\ |x| & \text{otherwise} \end{cases}$$
(28)

and to compare the certified value $c = E(H(\hat{p}))$ given in (12) against the Basharin, Hutcheson, and asymptotic approximations. The results obtained are shown in Fig. 2, where it can be seen that

for a previously selected number of possible patterns k and any of the underlying distributions here applied, the asymptotic estimate always provides the most accurate approximation (in terms of relative errors).

It is worth mentioning that we computed LRE and RE for $n \leq 1000$ since, as already mentioned, the formula for the exact expression $E(H(\hat{p}))$, given in Eq. (12), involves binomial coefficients that can only be computed with the aforementioned computational platforms when $n \leq 1000$. Nevertheless and despite the fact that these values of n are not large enough to be considered "asymptotic," the asymptotic estimate given in Eq. (25) is the most accurate

TABLE I. The largest *n* such that Mathematica v. 13.0.0 returns a value for Eq. (12) under two underlying distributions \mathcal{P} .

		i e		
720	120	24	6	\mathcal{P}
110	151	228	400	$\overline{\mathcal{P}_e}_{\mathcal{P}}$
	174	283	400 571	$\mathcal{P}_{e} \mathcal{P}_{L}$



FIG. 3. Histograms and boxplots of 10^6 values of the entropy from the \mathcal{P}_{L} model, $k \in \{6, 24, 120, 720\}$, and the series of length $n = 10^3$ k.

approximation of $\mathbb{E}(H(\hat{p}))$ in terms of LRE and RE for any combination of k, n, and underlying distribution even in those situations where $k \approx n$.

As previously mentioned, Eqs. (12) and (13) do not return usable values when implemented in either dependable numerical platforms or certified computer algebra systems. For instance, Table I shows the maximum *n* values for which the exact mean under \mathcal{P}_e and \mathcal{P}_L can be obtained using Mathematica¹⁷ in a computer Mac OS X ARM (64 bit). Notice that \mathcal{P}_e poses harder numerical problems than \mathcal{P}_L , but, in any case, these numerical limitations make the exact values from Eqs. (12) and (13) of little practical use; recall the rule-of-the-thumb $n \ge 100k$.

In order to study the behavior of expressions (12), (14), and (16) for long signals, in the following, we consider "certified" the value computed with independent simulations. First, we analyze the shape of the sample entropies from the \mathcal{P}_L model. For each value of $k \in \{6, 24, 120, 720\}$, we sample 10⁶ independent values of \hat{H} from the \mathcal{P}_L model and the series of length $n = 10^3$ k. Figure 3 shows the histograms and boxplots of these data. Table II shows the mean \hat{H} , median $q_{1/2}(\hat{H})$, standard deviation $s(\hat{H})$, asymmetry $\gamma_1(\hat{H})$, and excess kurtosis $\gamma_2(\hat{H})$ of the observed entropy over 10⁶ independent samples from the \mathcal{P}_L model, for four values of k = D!, along with approximate 95 % confidence intervals. It can be observed that, as k increases, so do the centrality measures of the samples (the mean \hat{H} and the median $q_{1/2}(\hat{H})$ coincide, suggesting no lack of symmetry). The dispersion, measured by the standard deviation $s(\hat{H})$, reduces when k increases. The asymmetry value, $\gamma_1(\hat{H})$, is slightly negative and reduces with increasing k, and the excess kurtosis $\gamma_2(\hat{H})$ is negligible in all cases.

Figure 4 shows the quantile–quantile plots of the entropies for the normal distribution. Notice that, although the histogram for k = 6 does not reveal any clear deviation from a normal density, the slight positive excess kurtosis explains the lack of fit of a few observations. Apart from that, the observations lie very close to the straight lines.

We now compare our "certified" values obtained by simulation with Hutcheson's exact values and third-order approximations.

TABLE II. Sample mean \hat{H} , median $q_{1/2}(\hat{H})$, standard deviation $s(\hat{H})$, asymmetry $\gamma_1(\hat{H})$, and excess kurtosis $\gamma_2(\hat{H})$ of the observed entropy over 10⁶ independent samples from the \mathcal{P}_{L} model, for four values of k = D!, along with approximate 95 % confidence intervals.

k	$\overline{\hat{H}}$	$q_{1/2}(\hat{H})$	$s(\hat{H})$	$\gamma_1(\hat{H})$	$\gamma_2(\hat{H})$
6	$0.9276 + 6.44 \times 10^{-6}$	$0.9276 + 8.09 \times 10^{-6}$	0.0033 +4.56 × 10 ⁻⁶	-0.0693 +4.80 × 10 ⁻³	0.0033 +9.60 × 10 ⁻³
24	$0.9451 \pm 1.97 \times 10^{-6}$	0.9451 $\pm 2.41 \times 10^{-6}$	$0.0010 \pm 1.40 \times 10^{-6}$	-0.0296 $\pm 4.80 \times 10^{-3}$	0.0004 $\pm 9.60 \times 10^{-3}$
120	0.9604 +5.92 × 10 ⁻⁷	0.9604 +7.34 × 10 ⁻⁷	0.0003 +4.19 × 10 ⁻⁷	-0.0115 +4.80 × 10 ⁻³	0.0020 +9.60 × 10 ⁻³
720	0.9707 $\pm 1.76 \times 10^{-7}$	0.9707 $\pm 2.18 \times 10^{-7}$		-0.0033 $\pm 4.80 \times 10^{-3}$	-0.0053 $\pm 9.60 \times 10^{-3}$



First, Fig. 5 shows, in log–log scales, the relationship between the relative error incurred by *H* to the mean of 10^6 independent samples $\overline{\hat{H}}$ as a function of the number of possible patterns *k* and of the factor that determines the number of observed patterns $n = \text{Factor} \times k$. For a sample size fixed, the relative error decreases with the number of possible patterns. When the number of possible patterns is fixed,

the relative error decreases as the sample size increases. The largest relative error is of the order of 5×10^{-3} ; it corresponds to the case k = 6 and n = 300 for which $H \approx 0.927$ 79 and $\overline{H} \approx 0.923$ 09. The smallest relative error occurs when k = 720and $n = 72 \times 10^6$. In such a case, it amounts to approximately 7.81987 $\times 10^{-6}$. Figure 6 shows, for each of the two extreme situations of minimum and maximum relative errors, the histogram of 10^6 observations, the underlying density function as a thin black line, and the asymptotic density distribution as a thick red line. We used a semilogarithmic scale.



FIG. 5. Relative error incurred by *H* to the mean of 10⁶ independent samples \hat{H} as a function of the number of possible patterns *k* and of the factor that determines the number of observed patterns $n = \text{Factor} \times k$.

It can be observed that the asymptotic standard deviation is acceptable for describing the data dispersion, and the asymptotic mean deviates from the observed mean, as Fig. 5 shows. Such a deviation is slight in the case shown in Fig. 6(a) and much more noticeable in Fig. 6(b). With this, we conclude that the relative error is an unreliable measure of the quality of a model.

Figures 5 and 6 suggest, thus, that there is room for an improved asymptotic model for the sample entropy \hat{H} . Our proposal for a corrected model stems from fusing the information we have about the asymptotic distribution of \hat{H} , namely, Eqs. (25) and (26), and the third-order corrected expected value of \hat{H} given in Eq. (16). With this, given the sequence $(\pi_1, \pi_2, \ldots, \pi_n)$ of symbols obtained from words of size D, first compute its histogram of proportions $\hat{p} = (\hat{p}_1, \hat{p}_2, \ldots, \hat{p}_k)$, where k = D! and $\hat{p}_\ell = \#\{j : \pi_j = \pi^{(\ell)}, 1 \le j \le n\}/n$, and then assume that $\hat{H} = -\sum_{\ell=1}^k \hat{p}_\ell \ln \hat{p}_\ell$ is an outcome



FIG. 6. Histograms of 10⁶ samples of \hat{H} under \mathcal{P}_{L} along with the empirical densities (thin black lines) and asymptotic models (thick red lines). (a) Case k = 6 and n = 300. (b) Case k = 720 and $n = 72 \times 10^6$.







FIG. 8. Maximum daily temperatures in Dublin, Edinburgh, and Miami, from 8 August 1992 until 30 December 2019, along with the histograms of k = 6 (left) and k = 24 (right) patterns.

from a random variable that has a normal distribution with mean

$$\mu_{n,p} = -\sum_{\ell=1}^{k} p_{\ell} \ln p_{\ell} - \frac{k-1}{2n} + \frac{1 - \sum_{\ell=1}^{k} p_{\ell}^{-1}}{12n^{2}} + \frac{\sum_{\ell=1}^{k} (p_{\ell}^{-1} - p_{\ell}^{-2})}{12n^{3}}$$
(29)

and variance given by

$$\sigma_{n,p}^{2} = \frac{1}{n} \sum_{\ell=1}^{k} p_{\ell} (1 - p_{\ell}) (\ln p_{\ell} + 1)^{2} - \frac{2}{n} \sum_{j=1}^{k-1} \sum_{\ell=j+1}^{k} p_{j} p_{\ell} (\ln p_{j} + 1) (\ln p_{\ell} + 1).$$
(30)

Figure 7 shows the histograms of 10⁶ samples of \hat{H} in two situations, namely, k = 6 and n = 300 [Fig. 7(a)], and k = 720 and $n = 72 \times 10^6$ [Fig. 7(b)]. In both cases, the samples were produced under $\mathcal{P}_{\rm L}$. The empirical densities are shown in thin black lines, while the corrected densities are shown in thick purple lines. The corrected model, i.e., the normal distribution with mean $\mu_{n,\hat{p}}$ and variance $\sigma_{n,\hat{p}}^2$, provides an excellent description of the observed data.

V. HYPOTHESIS TESTS AND APPLICATIONS

Let $\mathbf{x} = (x_1, x_2, \dots, x_{n_x})$ and $\mathbf{y} = (y_1, y_2, \dots, y_{n_y})$ be two independent time series of length $n_x = T_x + D_x - 1$ and $n_y = T_y + D_y - 1$, respectively. Let us also assume that these time series were generated by the stochastic (and maybe unknown) models \mathcal{M}_x and \mathcal{M}_y . We are interested in testing if these time series are statistically different. We choose to compare the entropy of their ordinal patterns among the many possible ways of performing such an assessment.

In this sense, we compute the series of symbols $\pi_1^x, \pi_2^x, \ldots, \pi_{D_x!}^y$ and $\pi_1^y, \pi_2^y, \ldots, \pi_{D_y!}^y$, and then, we find the histograms \hat{p}_x and \hat{p}_y . From Sec. III D, we know that Shannon's entropies $H(\hat{p}_x)$ and $H(\hat{p}_y)$ are random variables with asymptotic distributions $\mathcal{N}(\mu_{n_x,p_x}, \sigma_{n_x,p_x}^2)$ and $\mathcal{N}(\mu_{n_y,p_y}, \sigma_{n_y,p_y}^2)$, respectively, with means and variances given by Eqs. (29) and (30).

If **x** and **y** have the same underlying dynamics, then $H(\mathbf{p}_x) = H(\mathbf{p}_y)$, and we expect to observe $H(\hat{\mathbf{p}}_x) \approx H(\hat{\mathbf{p}}_y)$. Our test will then verify the following hypothesis:

$$\mathscr{H}_0: H(\boldsymbol{p}_{\mathbf{x}}) - H(\boldsymbol{p}_{\mathbf{y}}) = 0$$

vs the alternative

$$\mathscr{H}_1: H(\boldsymbol{p}_r) - H(\boldsymbol{p}_n) \neq 0.$$

It is, therefore, a bilateral test.

Our test statistic is

$$W = H(\hat{\boldsymbol{p}}_{x}) - H(\hat{\boldsymbol{p}}_{y}). \tag{31}$$

Using (29) and (30) and the assumption that the time series x and y are independent, it is straightforward that $W \xrightarrow{\mathscr{D}} \mathscr{N}(\mu_W, \sigma_W^2)$,

where $\mu_W = \mu_{n_x, p_x} - \mu_{n_y, p_y}$ and $\sigma_W^2 = \sigma_{n_x, p_x}^2 + \sigma_{n_y, p_y}^2$. Thus, for any observed $\eta > 0$, it holds that

$$\Pr\left(|H(\hat{\boldsymbol{p}}_{\boldsymbol{x}}) - H(\hat{\boldsymbol{p}}_{\boldsymbol{y}})| \le \eta \mid \mathscr{H}_{0}\right) \approx \Pr\left(|W| \le \eta \mid \mathscr{H}_{0}\right)$$
$$= 2\Phi\left(\frac{\eta - \mu_{W}}{\sigma_{W}}\right) - 1, \qquad (32)$$

where Φ is the standard Gaussian cumulative distribution function. Under \mathscr{H}_0 , the test statistic *W* is asymptotically distributed as

 $\mathcal{N}(0, \sigma_w^2)$, and therefore, the *p*-value of the observed test statistic is approximately $2(1 - \Phi(\varepsilon))$, where

$$\varepsilon = \frac{H(\hat{p}_x) - H(\hat{p}_y)}{\hat{\sigma}_W}$$
(33)

and

$$\hat{\sigma}_W = \sqrt{\sigma_{n_x,\hat{p}_x}^2 + \sigma_{n_y,\hat{p}_y}^2}.$$
(34)

Unilateral tests and other specific hypotheses can be easily obtained using that $H(\hat{p}_x) \sim \mathcal{N}(\mu_{n_x,\hat{p}_x}, \sigma_{n_x,p_x}^2)$ and that $H(\hat{p}_y) \sim \mathcal{N}(\mu_{n_y,\hat{p}_y}, \sigma_{n_y,p_y}^2)$. In particular, for the unilateral tests,

TABLE III. p-values of the hypothesis test applied to maximum temperature data.

Dublin vs Edinburgh					
Dublin	Edinburgh				
	k = 6	k = 24	k = 120	k = 720	
k = 6	0.0005	0.3294	0.6766	0.0830	
k = 24	$4.60 imes 10^{-5}$	0.0457	0.4694	0.6354	
k = 120	$1.48 imes10^{-6}$	0.0047	0.1124	0.6639	
k = 720	$4.98 imes 10^{-9}$	0.0001	0.0106	0.1594	

Dublin vs Edinburgh					
	Miami				
Dublin	k = 6	k = 24	k = 120	k = 720	
k = 6 $k = 24$ $k = 120$ $k = 720$	$\begin{array}{c} 8.78\times 10^{-8}\\ 4.85\times 10^{-3}\\ 1.48\times 10^{-1}\\ 8.66\times 10^{-1}\end{array}$	$\begin{array}{c} 1.68\times10^{-11}\\ 7.71\times10^{-6}\\ 1.54\times10^{-3}\\ 4.87\times10^{-5} \end{array}$	$\begin{array}{c} 3.33\times10^{-15}\\ 1.47\times10^{-8}\\ 1.26\times10^{-5}\\ 1.37\times10^{-3} \end{array}$	$0\\5.98 \times 10^{-12}\\2.89 \times 10^{-8}\\1.21 \times 10^{-5}$	

Dublin vs Edinburgh					
	Miami				
Edinburgh	k = 6	k = 24	k = 120	k = 720	
k = 6	0	0	0	0	
k = 24	$2.86 imes10^{-7}$	$9.05 imes 10^{-11}$	$4.33 imes 10^{-14}$	0	
k = 120	$7.86 imes10^{-4}$	$1.20 imes 10^{-6}$	$2.49 imes 10^{-9}$	1.25×10^{-12}	
k = 720	$5.83 imes 10^{-2}$	$4.57 imes 10^{-4}$	$3.30 imes 10^{-6}$	$7.13 imes 10^{-9}$	

(1) $\mathscr{H}_0: H(\boldsymbol{p}_x) - H(\boldsymbol{p}_y) \ge 0 \text{vs } \mathscr{H}_1: H(\boldsymbol{p}_x) - H(\boldsymbol{p}_y) < 0,$ (2) $\mathscr{H}_0: H(\boldsymbol{p}_x) - H(\boldsymbol{p}_y) \le 0 \text{vs } \mathscr{H}_1: H(\boldsymbol{p}_x) - H(\boldsymbol{p}_y) > 0,$

the *p*-values of the observed test statistics are approximately $\Phi(\varepsilon)$ and $1 - \Phi(\varepsilon)$, respectively, with ε defined in (33). These approximations follow the approach discussed by Boos and Stefanski (Sec. 3.6.1 in Ref. 18).

VI. CLIMATE DATA

Our goal in this analysis is to evaluate sequences that come from (i) the same underlying process and (ii) different dynamics. For this, we analyzed a series of maximum daily temperatures measured in three stations around the world: Dublin Phoenix Park (Ireland), Edinburgh Royal Botanic Garden (Scotland), and Miami International Airport (United States of America). We obtained these datasets from the Climate Data Online website, supported by the National Oceanic and Atmospheric Administration (NOAA) at https://www.ncei.noaa.gov/cdo-web/. They are part of the GHCN (Global Historical Climatology Network)-Daily data set, the world's most extensive collection of daily climatology measurements, that functions as the official archive for daily data from the Global Climate Observing System (GCOS) Surface Network (GSN). This data set contains observations of a variety of meteorological elements, including maximum and minimum temperatures, at more than 100 000 stations distributed across all continents.

We used daily data from 8 August 1992 until 30 December 2019. These daily observations are shown in Fig. 8 (in Fahrenheit degrees).

Dublin and Edinburgh have similar temperate oceanic climates, with few temperature differences. However, extreme temperatures might be a little below zero Celsius degrees ($32 \circ F$) a few days of the winter and make it into the mid-70s for some summer days. On the other hand, Miami has a tropical monsoon climate, with an average temperature of 77 ° F ($25 \circ C$) and extreme temperatures of 50 ° F in winter and 91.4 ° F in summer ($10 \circ C$ and $33 \circ C$, respectively). Figure 8 shows that, beyond these differences in the marginal properties, the series from Dublin and Edinburgh seem to follow a simple sine-cosine plus noise pattern closely. In contrast, the dynamics underlying the Miami data seem more complex: there is less variation except during January and February, where the maximum temperatures exhibit large deviations.

Figure 8 also shows the histograms of k = 6 (left) and k = 24 (right) patterns. Dublin and Edinburgh have similar histograms, while Miami's data have a predominant pattern. This configuration also appears when analyzing k = 120 and k = 720 patterns, but the visualization is too busy. Notice that it is impossible to make a visual comparison of histograms obtained with different embedding dimensions, although our test statistics allow for such an operation.

We applied the test introduced in Sec. V to contrast the null hypothesis between each possible pair of locations considering different values of the embedding dimension $D \in \{3, 4, 5, 6\}$ and, thus,



FIG. 9. Points in the $H \times C$ plane from (with boundaries in light gray) the three maximum daily temperature time series, along with the proposed confidence intervals for Shannon's entropy and different embedding dimensions *D*.

Chaos **32**, 113118 (2022); doi: 10.1063/5.0118706 Published under an exclusive license by AIP Publishing of the number of possible patterns $k \in \{6, 24, 120, 720\}$. In this case, all the time series have the same length n = 10000. The corresponding *p*-values are shown in Table III, from which we can see that H_0 is not rejected only for Dublin vs Edinburgh except for *k* taking larger values for Dublin than for Edinburgh, or k = 6 and k = 24 being the same value for both zones. These conclusions hold even after applying a conservative Bonferroni correction.¹⁹

We show in Fig. 9 the points in the $H \times C$ plane from the three maximum daily temperature time series, along with the proposed confidence intervals at the 0.1 % level for Shannon's entropy and different embedding dimensions $D \in \{3, 4, 5, 6\}$. The boundaries for each embedding dimension appear as light gray lines; see details about this kind of plot in the works by Chagas *et al.*⁸ and by Martin *et al.*²⁰ and the references therein. Notice that these are visual representations of the tests that contrast the entropies from series analyzed with the same embedding dimension; i.e., they depict the main diagonals of Table III.

Figure 9 shows that the maximum daily temperatures measured in Miami consistently have the smallest entropy and the largest complexity. The entropy carries enough information to discriminate its underlying dynamics from those that produce the measurements in Dublin and Edinburgh. Notice, also, that in this case, the complexity does not add information to the problem of discriminating the underlying processes that gave rise to the temperature measurements. The entropies from Dublin and Edinburgh are always very close, statistically indistinguishable at approximately the 95 % level of confidence in three (D = 4, 5, 6) out of four cases.

Whether the observed complexity is able or not to identify different underlying dynamics in Dublin and Edinburgh is an open question since, to date, we do not have expressions for their distribution.

VII. REMARKS AND CONCLUSIONS

Unlike the χ^2 test, the test statistic *W* we propose in (31) does not perform a bin-by-bin comparison. Our approach, on the one hand, represents an inevitable loss of information but, on the other hand, allows the comparison of entropies computed from different embedding dimensions.

The generality of our test promotes its application in federated learning, as it allows different sources to encode the data using different embedding.

Although we study Shannon's entropy from ordinal patterns, the approach to obtain the asymptotic distribution is valid for the Shannon entropy computed over transition graphs and their variants.

It is noteworthy that the presented derivations do not cover the extreme cases for which $H(\mathbf{p}) = 0$ or $H(\mathbf{p}) = 1$. The former does not involve any randomness. The latter was studied by Chagas *et al.*⁸ using an empirical approach.

ACKNOWLEDGMENTS

This work was partially funded by the National Council for Scientific and Technical Research—CONICET (Grant No. PIP2021 11220200103040CO), the São Paulo Research Foundation—FAPESP (Grant No. 2020/05121-4), and the National Council for Scientific and Technological Development—CNPq (Grant No. 312682/2021-2).

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflicts to disclose.

Author Contributions

E. T. C. Chagas: Conceptualization (equal); Data curation (equal); Formal analysis (equal); Funding acquisition (equal); Investigation (equal); Methodology (equal); Project administration (equal); Resources (equal); Software (equal); Supervision (equal); Validation (equal); Visualization (equal); Writing - original draft (equal); Writing - review & editing (equal). A. C. Frery: Conceptualization (equal); Data curation (equal); Formal analysis (equal); Funding acquisition (equal); Investigation (equal); Methodology (equal); Project administration (equal); Resources (equal); Software (equal); Supervision (equal); Validation (equal); Visualization (equal); Writing – original draft (equal); Writing – review & editing (equal). J. Gambini: Conceptualization (equal); Data curation (equal); Formal analysis (equal); Funding acquisition (equal); Investigation (equal); Methodology (equal); Project administration (equal); Resources (equal); Software (equal); Supervision (equal); Validation (equal); Visualization (equal); Writing - original draft (equal); Writing - review & editing (equal). M. M. Lucini: Conceptualization (equal); Data curation (equal); Formal analysis (equal); Funding acquisition (equal); Investigation (equal); Methodology (equal); Project administration (equal); Resources (equal); Software (equal); Supervision (equal); Validation (equal); Visualization (equal); Writing - original draft (equal); Writing - review & editing (equal). H. S. Ramos: Conceptualization (equal); Data curation (equal); Formal analysis (equal); Funding acquisition (equal); Investigation (equal); Methodology (equal); Project administration (equal); Resources (equal); Software (equal); Supervision (equal); Validation (equal); Visualization (equal); Writing - original draft (equal); Writing - review & editing (equal). A. A. Rey: Conceptualization (equal); Data curation (equal); Formal analysis (equal); Funding acquisition (equal); Investigation (equal); Methodology (equal); Project administration (equal); Resources (equal); Software (equal); Supervision (equal); Validation (equal); Visualization (equal); Writing - original draft (equal); Writing - review & editing (equal).

DATA AVAILABILITY

This work employed simulated data in Sec. IV. The temperature datasets analyzed in Sec. VI are openly available at https://www.ncei.noaa.gov/cdo-web/.

REFERENCES

¹C. Bandt and B. Pompe, "Permutation entropy: A natural complexity measure for time series," Phys. Rev. Lett. **88**, 174102-1–174102-4 (2002).

²O. A. Rosso, H. A. Larrondo, M. T. Martín, A. Plastino, and M. A. Fuentes, "Distinguishing noise from chaos," Phys. Rev. Lett. **99**, 154102 (2007).

³M. Zanin, L. Zunino, O. A. Rosso, and D. Papo, "Permutation entropy and its main biomedical and econophysics applications: A review," Entropy 14, 1553–1577 (2012).

⁴J. Chen, C. Lin, D. Peng, and H. Ge, "Fault diagnosis of rotating machinery: A review and bibliometric analysis," IEEE Access **8**, 224985–225003 (2020).

⁵E. Chagas, A. C. Frery, O. A. Rosso, and H. S. Ramos, "Analysis and classification of SAR textures using information theory," IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens. **14**, 663–675 (2021).

⁶L. Zunino and H. Ribeiro, "Discriminating image textures with the multiscale two-dimensional complexity-entropy causality plane," Chaos, Solitons Fractals **91**, 679–688 (2016).

⁷H. V. Ribeiro, L. Zunino, E. K. Lenzi, P. A. Santoro, and R. S. Mendes, "Complexity-entropy causality plane as a complexity measure for twodimensional patterns," PLoS One 7, e40689 (2012).

⁸E. T. C. Chagas, M. Queiroz-Oliveira, O. A. Rosso, H. S. Ramos, C. G. S. Freitas, and A. C. Frery, "White noise test from ordinal patterns in the entropy-complexity plane," Int. Stat. Rev. **90**, 374 (2022).

⁹G. Basharin, "On a statistical estimate for the entropy of a sequence of independent random variables," Theory Probab. Its Appl. 4, 333–336 (1959).

¹⁰K. Hutcheson, "A test for comparing diversities based on the Shannon formula," J. Theor. Biol. **29**, 151–154 (1970). ¹¹K. Hutcheson and L. R. Shenton, "Some moments of an estimate of Shannon's measure of information," Commun. Stat.-Theory Methods **3**, 89–94 (1974).

¹²N. L. Johnson, S. Kotz, and A. W. Kemp, "Univariate discrete distributions," in *Wiley Series in Probability and Mathematical Statistics*, 2nd ed. (John Wiley & Sons, New York, 1993).

¹³G. Casella and R. L. Berger, *Statistical Inference*, 2nd ed. (Duxbury, Pacific Grove, CA, 2002).

¹⁴P. Mukhopadhyay, *Complex Surveys: Analysis of Categorical Data* (Springer, 2016).

¹⁵E. L. Lehmann and G. Casella, *Theory of Point Estimation* (Springer Science & Business Media, 2006).

¹⁶M. Almiron, B. L. Vieira, A. L. C. Oliveira, A. C. Medeiros, and A. C. Frery, "On the numerical accuracy of spreadsheets," J. Stat. Softw. 34, 1–29 (2010).

¹⁷Mathematica version 13.0.0, Wolfram Research, Inc., Champaign, IL, 2021.

¹⁸D. D. Boos and L. A. Stefanski, *Essential Statistical Inference: Theory and Methods*, Springer Texts in Statistics (Springer, New York, 2014).

¹⁹S. Nakagawa and I. C. Cuthill, "Effect size, confidence interval and statistical significance: A practical guide for biologists," Biol. Rev. 82, 591–605 (2007).
²⁰M. T. Martin, A. Plastino, and O. A. Rosso, "Generalized statistical complex-

²⁰M. I. Martin, A. Plastino, and O. A. Rosso, "Generalized statistical complexity measures: Geometrical and analytical properties," Physica A **369**, 439–462 (2006).